Patient's Name: _	
	Michael A. Ruchim. <i>I</i>

Michael A. Ruchim, M.D. 680 N. Lake Shore Dr. – Suite 118 Chicago, IL 60611 (312) 503-6000

# **MORNING COLONOSCOPY INSTRUCTIONS**

DATE OF TEST:	TIME:	a.m. /	/ p.m.
(A	ARRIVE 30-45 MINUTES BEFORE YOUR SCHEDULED APPOINTME	NT)	

LOCATION: Northwestern Memorial Hospital G.I. Lab – 4<sup>th</sup> Floor Galter Pavilion

675 N. St. Clair - Chicago, IL 60611

## IN PREPARATION FOR THE PROCEDURE

- ➤ Have the (enclosed) prescription for MoviPrep® filled.
- Fill out the GI LAB QUESTIONNAIRE and PATIENT MEDICATION LIST (enclosed).
- Do not take **PEPTO BISMOL** or **IRON TABLETS** for **7 days** (one week) prior to the exam.
- Do not take any ASPIRIN, PLAVIX or any anti-inflammatory medications, such as MOTRIN, ALEVE, ADVIL or IBUPROFIN for 5 days prior to the procedure.
- If you take **COUMADIN/WARFARIN**, contact your Primary Care Physician so he/she can advise you when to stop taking it prior to the test.
- If you are on INSULIN, contact your Primary Care Physician regarding dosage adjustment.
- All medications **NOT LISTED ABOVE** may be taken normally as directed.

#### THE DAY BEFORE THE PROCEDURE

- In the morning, prepare one liter of MoviPrep® according to the instructions on the package and refrigerate.
- You must follow a clear liquid diet the day before the test. It is important to drink at least one glass of a clear liquid EVERY HOUR throughout the day.

## YOU *MAY* HAVE:

- ✓ Apple Juice
- ✓ Cranberry Juice
- ✓ Black Coffee
- ✓ Tea
- ✓ White Grape Juice

- ✓ Chicken or Beef Broth
- ✓ Popsicles®
- ✓ Jell-O® (NOT RED)
- ✓ Gatorade
- ✓ Soda Pop

#### YOU MAY NOT HAVE:

- ✓ Sherbet
- ✓ Ice Cream

- ✓ Milk or Milk Products
- ✓ Orange Juice

(CONTINUED over)

## THE DAY BEFORE THE PROCEDURE (continued)

- ➤ Between 6:00 and 7:00 p.m., drink the MoviPrep® as instructed; one full glass (8 oz.) every 10-15 minutes. Drink each glass quickly rather than drinking small amounts continuously. *BE SURE TO DRINK ALL OF THE SOLUTION.*
- $\triangleright$  When finished, prepare the 2<sup>nd</sup> liter of MoviPrep® and refrigerate.
- Approximately 4-5 hours before the colonoscopy, drink the 2<sup>nd</sup> liter of MoviPrep® as the 1<sup>st</sup>; one full glass (8 oz.) every 10-15 minutes. This could require waking up during the night.
- Additionally, drink 4 glasses (32 oz.) of water over the course of the evening.
- ➤ You must not have ANYTHING BY MOUTH after the final dose of MoviPrep®

## THE DAY OF THE PROCEDURE.

Bring the completed **GI LAB QUESTIONNAIRE** and **PATIENT MEDICATION LIST** enclosed with you to registration. Report to the hospital approximately 30-45 minutes prior to your appointment time.

You <u>MUST</u> be accompanied by a friend or relative to drive and/or assist you home. Walking, taxi or public transportation is not allowed unless you have another adult with you. This is strict hospital policy and <u>failure to comply will result in cancellation of the procedure</u>. If you are unable to arrange for an adult to accompany you, you may call the hospital at (312) 926-7614 to discuss transportation options. This MUST be coordinated prior to the date of the test.

\*\*\*If you must cancel or reschedule your appointment, please give 72-HOUR NOTICE to Dr. Ruchim's office at (312) 503-6000.\*\*\*

If you have any questions or are experiencing any difficulty with the preparation, contact Dr. Ruchim at (312) 503-6000.

Please contact Northwestern Patient Accounts at (312) 926-3642 if you have any billing or insurance questions.